

CONSTRUCTING EXACT LAGRANGIAN IMMERSIONS WITH FEW DOUBLE POINTS

TOBIAS EKHOLM, YAKOV ELIASHBERG, EMMY MURPHY, AND IVAN SMITH

ABSTRACT. We establish, as an application of the results from [13], an h -principle for exact Lagrangian immersions with transverse self-intersections and the minimal, or near-minimal number of double points. One corollary of our result is that any orientable closed 3-manifold admits an exact Lagrangian immersion into standard symplectic 6-space \mathbb{R}_{st}^6 with exactly one transverse double point. Our construction also yields a Lagrangian embedding $S^1 \times S^2 \rightarrow \mathbb{R}_{\text{st}}^6$ with vanishing Maslov class.

1. INTRODUCTION

Lagrangian self-intersections. In this paper we study the problem of constructing Lagrangian immersions with the minimal possible number of transverse self-intersection points. It is well known that the existence of a Lagrangian embedding imposes strong topological constraints (e.g. Gromov's theorem about non-existence of exact Lagrangian submanifolds in standard symplectic $2n$ -space, $\mathbb{R}_{\text{st}}^{2n} = (\mathbb{R}^{2n}, \sum_{i=1}^n dx_i \wedge dy_i)$), and also that, in many cases, two Lagrangian submanifolds must intersect in more points than is suggested by topological intersection theory alone (e.g. results confirming Arnold's conjectures). In view of this, it was expected that there should be similar lower bounds on the minimal number of double points for Lagrangian immersions, e.g. that an exact Lagrangian immersion of an n -torus into $\mathbb{R}_{\text{st}}^{2n}$ with transverse self-intersections would have at least 2^{n-1} double points.

Bounds of this kind have been proved for Lagrangian immersions satisfying additional conditions. For instance, it was shown in [6, 8] that any self-transverse exact Lagrangian immersion $f: L \rightarrow \mathbb{R}_{\text{st}}^{2n}$ of a closed n -manifold, for which the Legendrian lift $\tilde{f}: L \rightarrow \mathbb{R}_{\text{st}}^{2n} \times \mathbb{R}$ has Legendrian homology algebra that admits an augmentation, satisfies the following analogue of the Morse inequalities: the number of double points of f is at least $\frac{1}{2} \sum_{j=0}^n \text{rank}(H_j(L))$.

Date: March 2013.

T.E. is partially supported by Swedish Research Council Grant 2012-2365 and by the Knut and Alice Wallenberg Foundation as a Wallenberg Scholar.

Y.E. is partially supported by NSF grant DMS-1205349.

I.S. is partially supported by grant ERC-2007-StG-205349 from the European Research Council.

Here we prove a surprising result in sharp contrast to such lower bounds: if no extra constraints are imposed then Lagrangian immersions into symplectic manifolds of dimension > 4 with *nearly the minimal* number of self-intersection points satisfy a certain h -principle. Let us introduce the notation needed to state the result. Let X be an oriented $2n$ -manifold and $f: L \rightarrow X$ a proper smooth immersion of a connected n -manifold. If L is non-compact we will assume that f is an embedding outside of a compact set. Following Whitney [22], if f is self-transverse we assign a self-intersection number $I(f)$ to f , where $I(f)$ is the mod 2 number of self-intersection points if n is odd or L is non-orientable, and the algebraic number of self-intersection points counted with their intersection signs if n is even and L is orientable. Note that $I(f)$ depends only on the regular homotopy class of f provided that L is closed or that the regular homotopy is an isotopy at infinity. When $n > 2$ and X is simply connected, a theorem of Whitney [22] asserts that an immersion f is regularly homotopic to an embedding if and only if $I(f) = 0$. If $f: L \rightarrow X$ is a self-transverse immersion, then we let $\text{SI}(f) \geq 0$ denote its total number of double points. Clearly, $\text{SI}(f) \geq |I(f)|$.

Similarly, given an orientable $(2n-1)$ -dimensional manifold Y and an $(n-1)$ -dimensional manifold Λ , consider a smooth regular homotopy $h_t: \Lambda \rightarrow Y$, $0 \leq t \leq 1$, which connects embeddings h_0 and h_1 and is an isotopy outside of a compact set. Let $H: \Lambda \times [0, 1] \rightarrow Y \times [0, 1]$ be given by $H(x, t) = (h_t(x), t)$; then H is an immersion. We say that the regular homotopy h_t has transverse intersections if the immersion H has transverse double points. In this case we define $I(\{h_t\}_{t \in [0, 1]}) := I(H)$ and $\text{SI}(\{h_t\}_{t \in [0, 1]}) := \text{SI}(H)$. Note that if n is even and Λ is orientable then $I(\{h_t\}_{t \in [0, 1]}) = -I(\{h_{1-t}\}_{t \in [0, 1]})$.

Consider next a Lagrangian regular homotopy, $f_t: L \rightarrow X$, $0 \leq t \leq 1$, and write $F: L \times [0, 1] \rightarrow X$ for $F(x, t) = f_t(x)$. Let α denote the 1-form on $L \times [0, 1]$ defined by the equation $\alpha := \iota_{\partial/\partial t}(F^*\omega)$, where ι denotes contraction and t is the coordinate on the second factor of $L \times [0, 1]$. Then the restrictions $\alpha_t := \alpha|_{L \times \{t\}}$ are closed for all t . We call the Lagrangian regular homotopy f_t a *Hamiltonian regular homotopy* if the cohomology class $[\alpha_t] \in H^1(L)$ is independent of t . The following theorem is our main result:

THEOREM 1.1. *Suppose that X is a simply connected $2n$ -dimensional symplectic manifold, $n > 2$. If $f: L \rightarrow X$ is a Lagrangian immersion then there exists a Hamiltonian regular homotopy $f_t: L \rightarrow X$, $0 \leq t \leq 1$, with $f_0 = f$ such that f_1 is self-transverse and*

$$\text{SI}(f_1) = \begin{cases} 1, & \text{if } n \text{ is odd or } L \text{ is non-orientable and } I(f_0) = 1; \\ 2, & \text{if } n \text{ is odd or } L \text{ is non-orientable and } I(f_0) = 0; \\ |I(f_0)|, & \text{if } n \text{ is even, } L \text{ is orientable and } (-1)^{\frac{n}{2}} I(f_0) < 0; \\ |I(f_0)| + 2, & \text{if } n \text{ is even, } L \text{ is orientable and } (-1)^{\frac{n}{2}} I(f_0) \geq 0. \end{cases}$$

REMARK 1.2. There is a version of Theorem 1.1 in the non-simply connected case where the intersection number $I(f_0)$ is defined as an element of the group ring of $\pi_1(X)$ and where $|I(f_0)|$ denotes an appropriate norm on this ring. For simplicity, we focus on the simply connected case in this paper.

Theorem 1.1 is proved in Section 4.1 as an application of results in [13].

This result has the following consequences for exact Lagrangian immersions into $\mathbb{R}_{\text{st}}^{2n}$. Let L be an n -dimensional closed manifold. Recall that according to Gromov's h -principle for Lagrangian immersions the triviality of the complexified tangent bundle $TL \otimes \mathbb{C}$ is a necessary and sufficient condition for the existence of an exact Lagrangian immersion $L \rightarrow \mathbb{R}_{\text{st}}^{2n}$, while exact Lagrangian regular homotopy classes are in natural one to one correspondence with homotopy classes of trivializations of $TL \otimes \mathbb{C}$. We write $s(L)$ for the minimal number of double points of a self-transverse exact Lagrangian immersion $L \rightarrow \mathbb{R}_{\text{st}}^{2n}$. Given a homotopy class σ of trivializations of $TL \otimes \mathbb{C}$, the refined invariant $s(L, \sigma)$ denotes the minimal number of double points of an exact self-transverse Lagrangian immersion $L \rightarrow \mathbb{R}_{\text{st}}^{2n}$ representing the exact Lagrangian regular homotopy class σ .

COROLLARY 1.3. *Let L be an n -dimensional closed manifold with $TL \otimes \mathbb{C}$ trivial and let σ be a homotopy class of trivializations of $TL \otimes \mathbb{C}$. Then the following hold:*

- (1) *If $n > 1$ is odd or if L is non-orientable, then $s(L, \sigma) \in \{1, 2\}$.*
- (2) *If $n = 1$, then $s(L) = 1$ and there exist σ with $s(L, \sigma) = d$ for any integer $d > 0$; if $n = 3$, then $s(L) = 1$ and for one of the two regular homotopy classes σ , $s(L, \sigma) = 2$.*
- (3) *If n is even and L is orientable, then for $\chi(L) < 0$, $s(L, \sigma) = \frac{1}{2}|\chi(L)|$, and for $\chi(L) \geq 0$, either $s(L, \sigma) = \frac{1}{2}\chi(L)$ or $s(L, \sigma) = \frac{1}{2}\chi(L) + 2$.*

Corollary 1.3 is proved in Section 4.2; Section 4.3 then give more detailed information about the case when n is odd, respectively discuss the Lagrangian embeddings obtained from these immersions by Lagrange surgery. The case $n = 2$ in both (1) and (3) above does not follow from our proof of Theorem 1.1; rather, these are results of Sauvaget [20], who gave direct geometric constructions of self-transverse exact Lagrangian immersions of both oriented and non-oriented surfaces. In particular, Sauvaget constructed, as the key point for his result, an exact immersed genus two surface in \mathbb{C}^2 with exactly one double point. In Appendix A we describe a higher dimensional analogue of that construction.

It is interesting to compare Corollary 1.3 with the results of [9, 10] which show that any homotopy n -sphere Σ that admits a Lagrangian immersion into $\mathbb{R}_{\text{st}}^{2n}$ with exactly one transverse double point of even Maslov grading and with induced trivialization of the complexified tangent bundle homotopic to that of the Whitney immersion of the standard n -sphere¹, must bound a parallelizable $(n + 1)$ -manifold. If n is even, both the Maslov grading condition and the homotopy condition are automatically satisfied, and moreover the standard n -sphere is the only homotopy n -sphere that bounds a parallelizable $(n + 1)$ -manifold. Thus, if n is even the standard n -sphere is the only homotopy n -sphere that admits a self-transverse Lagrangian immersion into Euclidean space with only one double point. This means in particular that in the case when $\dim(L)$ is even and $\chi(L) > 0$, $s(L)$

¹i.e. the trivializations $\sigma: TS^n \otimes \mathbb{C} \rightarrow \mathbb{C}^n$ and $\tilde{\sigma}: T\Sigma \otimes \mathbb{C} \rightarrow \mathbb{C}^n$ are related as $\tilde{\sigma} = \sigma \circ (h \otimes \mathbb{C})$, where $h: T\Sigma \rightarrow TS$ is a bundle isomorphism covering a homotopy equivalence $\Sigma \rightarrow S^n$.

is generally not determined by the homotopy type of L . The following result constrains the homotopy type of a manifold for which this phenomenon may occur.

THEOREM 1.4. *Let L be an even dimensional spin manifold with $\chi(L) > 0$. If $s(L) = \frac{1}{2}\chi(L)$ then $\pi_1(L) = 1$ and $H_{2k+1}(L) = 0$ for all k . In particular if $\dim L > 4$ then L has the homotopy type of a CW-complex with $\chi(L)$ even-dimensional cells and no odd-dimensional cells.*

Theorem 1.4 is proved in Section 4.4. The proof uses lifted linearized Legendrian homology, introduced in [10] following ideas in [5]. Note that this result can be viewed as an obstruction to an h -principle for exact Lagrangian immersions having the minimal, rather than near-minimal, number of self-intersection points.

Surgery, Lagrangian embeddings and the Maslov class. In [19] Polterovich describes a local Lagrangian surgery construction which resolves a double point of a Lagrangian immersion. Let Q_+ denote the manifold $S^1 \times S^{n-1}$, and Q_- the mapping torus of an orientation-reversing involution of S^{n-1} . Given a Lagrangian immersion $f: L \rightarrow X$ of an oriented n -manifold with a single double point p , [19, Propositions 1 & 2] imply:

- (1) if n is odd, there are Lagrangian embeddings $L \# Q_{\pm} \rightarrow X$;
- (2) if n is even, there is a Lagrangian embedding $L \# Q_{\epsilon} \rightarrow X$, where the sign of ϵ is given by $(-1)^{n(n-1)/2+1} \text{sign}(p)$,

where $\text{sign}(p) \in \{\pm 1\}$ denotes the intersection index of the double point. Combining this with Corollary 1.3 yields:

COROLLARY 1.5. *Let Y be a closed orientable 3-manifold. Then there is a Lagrangian embedding $Y \# (S^1 \times S^2) \rightarrow \mathbb{R}_{\text{st}}^6$.*

The question of determining the minimal number k for which there is a Lagrangian embedding $Y \# k(S^1 \times S^2) \rightarrow \mathbb{R}_{\text{st}}^6$ was raised explicitly by Polterovich [19, Remark 2].

In another direction, in each odd dimension $n = 2k + 1$ our construction yields a Lagrangian immersion of the sphere $S^n \rightarrow \mathbb{R}_{\text{st}}^{2n}$ with a single double point of Maslov grading 1 (note the double point of the Whitney sphere has Maslov grading n). Using Lagrange surgery we conclude:

COROLLARY 1.6. *There exists a Lagrangian embedding $S^1 \times S^{2k} \rightarrow \mathbb{R}_{\text{st}}^{4k+2}$ for which the generator of the first homology of positive action has non-positive Maslov index $2 - 2k$. In particular, there exists a Lagrangian embedding $S^1 \times S^2 \rightarrow \mathbb{R}_{\text{st}}^6$ with zero Maslov class.*

The (non-)existence question for Lagrangian embeddings into $\mathbb{R}_{\text{st}}^{2n}$ with vanishing Maslov class is a well-known problem in symplectic topology. Viterbo proved in [21] that if L

admits a metric of non-positive sectional curvature then any Lagrangian embedding $L \rightarrow \mathbb{R}_{\text{st}}^{2n}$ has non-trivial Maslov class, whilst Fukaya, Oh, Ohta and Ono infer the same conclusion in Theorem K of [15] whenever L is a spin manifold with $H^2(L; \mathbb{Q}) = 0$. Corollary 1.6 shows that the hypotheses in these theorems play more than a technical role; in particular, the assumption on second cohomology in the latter result cannot be removed. We point out that the Lagrangian embedding in Corollary 1.6 is not monotone. In the monotone case, the Maslov index of the generator of the first homology of positive action necessarily equals 2, see [14, Proposition 2.10] (and references therein) and [5, Theorem 1.5 (b)].

Plan of the paper. The paper is organized as follows. In Section 2 we recall the notions and the main results of the theory of loose Legendrian knots from [18]. In Section 3 we formulate the h -principle for Lagrangian embeddings with conical singularities, established in [13], and deduce from it its own generalization: in the presence of a conical singularity, Lagrangian immersions with the minimal number of self-intersections abide an h -principle. Theorem 1.1 is then proved in Section 4.1 as an application of this h -principle. Corollary 1.3 and related explicit results about Lagrangian immersions into $\mathbb{R}_{\text{st}}^{2n}$ are proved in Sections 4.2, 4.3, and 4.4. As is typical with h -principles, Theorem 1.1 does not provide explicit constructions of Lagrangian immersions with the minimal number of double points. In Appendix A we complement Theorem 1.1 by constructing an explicit exact Lagrangian immersion of $P = (S^1 \times S^{n-1}) \# (S^1 \times S^{n-1})$ into $\mathbb{R}_{\text{st}}^{2n}$ with exactly one transverse double point (in particular, yielding an immersion in each dimension violating the Arnold-type bound which pertains in the presence of linearizable Legendrian homology algebra). Our construction is a generalization of Sauvaget's construction in the case $n = 2$. The construction of the Appendix seems to have no elementary relation to the arguments earlier in the paper; in general, our immersions with few double points are obtained by first introducing many double points and then cancelling them in pairs, whilst in the Appendix considerable effort is expended to keep the Lagrangians swept by appropriate Legendrian isotopies embedded.

Acknowledgements. T.E. and I.S. are grateful to François Laudenbach for drawing their attention to Sauvaget's work. Y.E. is grateful to the Simons Center for Geometry and Physics where a part of this paper was completed.

2. LOOSE LEGENDRIAN KNOTS

The theory of loose Legendrian knots and the h -principle for Lagrangian caps with loose Legendrian ends, developed in [18] and [13], respectively, are crucial for the proof of our main result, Theorem 1.1. In this section we recall the concepts and results from these theories that will be used in later sections.

2.1. Stabilization. We start with a discussion of the *stabilization construction* for Legendrian submanifolds, see [11, 4, 18].

Consider standard contact \mathbb{R}^{2n-1} :

$$\mathbb{R}_{\text{st}}^{2n-1} = \left(\mathbb{R}^{2n-1}, \xi_{\text{st}} = \ker \left(dz - \sum_1^{n-1} y_i dx_i \right) \right),$$

where $(x_1, y_1, \dots, x_{n-1}, y_{n-1}, z)$ are coordinates in \mathbb{R}^{2n-1} , with the Legendrian coordinate subspace $\Lambda_0 \subset \mathbb{R}_{\text{st}}^{2n-1}$ given by

$$\Lambda_0 = \{(x_1, y_1, \dots, x_{n-1}, y_{n-1}, z) : x_1 = y_2 = y_3 = \dots = y_{n-1} = 0, z = 0\}.$$

Then $(\mathbb{R}_{\text{st}}^{2n-1}, \Lambda_0)$ is a local model for any Legendrian submanifold in a contact manifold. More precisely, if $\Lambda \subset Y$ is any Legendrian $(n-1)$ -submanifold of a contact $(2n-1)$ -manifold Y then any point $p \in \Lambda$ has a neighborhood $\Omega \subset Y$ that admits a map

$$\Phi : (\Omega, \Lambda \cap \Omega) \rightarrow (\mathbb{R}_{\text{st}}^{2n-1}, \Lambda_0), \quad \Phi(p) = 0,$$

which is a contactomorphism onto a neighborhood of the origin.

We will carry out the stabilization construction in a local model that is slightly different from $(\mathbb{R}_{\text{st}}^{2n-1}, \Lambda_0)$, which we discuss next. Let $F : \mathbb{R}_{\text{st}}^{2n-1} \rightarrow \mathbb{R}_{\text{st}}^{2n-1}$ denote the contactomorphism,

$$F(x_1, y_1, \dots, x_{n-1}, y_{n-1}, z) = \left(x_1 + \frac{1}{2}y_1^2, y_1, x_2, y_2, \dots, x_{n-1}, y_{n-1}, z + \frac{1}{3}y_1^3 \right).$$

Then F maps Λ_0 to

$$\Lambda_{\text{cu}} = \left\{ (x_1, y_1, \dots, x_{n-1}, y_{n-1}, z) : x_1 = \frac{1}{2}y_1^2, y_2 = \dots = y_{n-1} = 0, z = \frac{1}{3}y_1^3 \right\}$$

In the language of Appendix A.2, the front Γ_{cu} of Λ_{cu} in $\mathbb{R}^{n-1} \times \mathbb{R}$ is the product of $\mathbb{R}^{n-2} \subset \mathbb{R}^{n-1}$ and the standard cusp $\{(x_1, z) : 9z^2 = 8x_1^3\}$ in $\mathbb{R} \times \mathbb{R}$. In particular, the two branches of the front are graphs of the functions $\pm h$, where

$$h(x) = h(x_1, \dots, x_{n-1}) = \frac{2\sqrt{2}}{3}x_1^{\frac{3}{2}},$$

defined on the half-space $\mathbb{R}_+^{n-1} := \{x = (x_1, \dots, x_{n-1}) : x_1 \geq 0\}$.

Let U be a domain with smooth boundary contained in the interior of \mathbb{R}_+^{n-1} , $U \subset \text{int}(\mathbb{R}_+^{n-1})$. Pick a non-negative function $\phi : \mathbb{R}_+^{n-1} \rightarrow \mathbb{R}$ with the following properties: ϕ has compact support in $\text{int}(\mathbb{R}_+^{n-1})$, the function $\tilde{\phi}(x) := \phi(x) - 2h(x)$ is Morse, $U = \tilde{\phi}^{-1}([0, \infty))$, and 0 is a regular value of $\tilde{\phi}$. Consider the front Γ_{cu}^U in $\mathbb{R}^{n-1} \times \mathbb{R}$ obtained from Γ_{cu} by replacing the lower branch of Γ_{cu} , i.e. the graph $z = -h(x)$, by the graph $z = \phi(x) - h(x)$. Since ϕ has compact support, the front Γ_{cu}^U coincides with Γ_{cu} outside a compact set. Consequently, the Legendrian embedding $\Lambda_{\text{cu}}^U : \mathbb{R}^{n-1} \rightarrow \mathbb{R}^{2n-1}$ defined by the front Γ_{cu}^U coincides with Λ_{cu} outside a compact set.

LEMMA 2.1 ([4, 18]). *There exists a compactly supported Legendrian regular homotopy $\Lambda_{\text{cu}; t}$, $t \in [0, 1]$ connecting Λ_{cu} to Λ_{cu}^U with $\text{SI}(\{\Lambda_{\text{cu}; t}\}_{t \in [0, 1]})$ equal to the number of critical points of $\phi(x) - h(x)$, and with $I(\{\Lambda_{\text{cu}; t}\}_{t \in [0, 1]}) = (-1)^{k-1}\chi(U)$ if $n = 2k$ and $I(\{\Lambda_{\text{cu}; t}\}_{t \in [0, 1]}) \equiv \chi(U) \pmod{2}$ if n is odd.*

Proof. It is straightforward to construct a family of functions $\phi_t(x)$, $t \in [0, 1]$, with the following properties:

- (1) $\phi_0(x) = 0$ and $\phi_1(x) = \phi(x)$;
- (2) the functions are monotonically increasing in t ;
- (3) for each critical point x of $\tilde{\phi}|_U = (\phi - 2h)|_U$, there is a unique $t \in [0, 1]$ such that x is a critical point of the function $\phi_t - 2h$, of the same index and of critical value 0.

We associate with ϕ_t the front $\Gamma_{\text{cu};t}$ obtained from $\Gamma_{\text{cu}} = \Gamma_{\text{cu};0}$ by replacing the lower branch of Γ_{cu} , which by definition is the graph $z = -h(x)$, by the graph $z = \phi_t(x) - h(x)$. The Legendrian regular homotopy $\Lambda_{\text{cu};t}$ determined by this family of fronts has transverse self-intersections which correspond to critical points of $\phi_t - 2h$ of critical value 0. One can show that when $n = 2k$ the sign of each intersection point equals $(-1)^{\text{ind } x + k - 1}$, where $\text{ind } x$ is the Morse index of the critical point x of the function $(\phi - 2h)|_U$, see [4]. \square

To transport the stabilization construction to our standard local model, let $\Lambda_0^U = F^{-1}(\Lambda_{\text{cu}}^U)$ and $\Lambda_{0;t} = F^{-1}(\Lambda_{\text{cu};t})$, where $\Lambda_{\text{cu};t}$, $t \in [0, 1]$, is the regular Legendrian homotopy constructed in Lemma 2.1. Then $\Lambda_{0;t}$, $t \in [0, 1]$, is a compactly supported Legendrian regular homotopy connecting Λ_0 to Λ_0^U .

Consider a Legendrian $(n - 1)$ -submanifold Λ of a contact $(2n - 1)$ -manifold Y and a point $p \in \Lambda$. Fix a neighborhood $\Omega \subset Y$ of p and a contactomorphism

$$\Phi: (\Omega, \Lambda \cap \Omega) \rightarrow (\mathbb{R}_{\text{st}}^{2n-1}, \Lambda_0).$$

Replacing $\Omega \cap \Lambda$ with $\Phi^{-1}(\Lambda_0^U)$ we get a Legendrian embedding Λ^U which coincides with Λ outside of Ω , and replacing it with $\Phi^{-1}(\Lambda_{0;t})$ we get a Legendrian regular homotopy Λ_t , $t \in [0, 1]$ connecting Λ to Λ^U with $\text{SI}(\{\Lambda_t\}_{t \in [0,1]})$ equal to the minimal number of critical points of a Morse function on U which attains its minimum value on ∂U , and with $I(\{\Lambda_t\}_{t \in [0,1]}) = (-1)^{k-1} \chi(U)$ if $n = 2k$ and $I(\{\Lambda_t\}_{t \in [0,1]}) \equiv \chi(U) \pmod{2}$ if n is odd. We say that Λ^U is obtained from U via U -stabilization in Ω . The most important case for us will be when U is the ball. We say in this case that Λ^U is the *stabilization* of Λ in Ω or simply the stabilization of Λ .

Let ξ denote the contact plane field on Y , and note that there is an induced field of (conformally) symplectic 2-forms on ξ . We say that two Legendrian embeddings $f_0, f_1: \Lambda \rightarrow Y$ are *formally Legendrian isotopic* if there exists a smooth isotopy $f_t: \Lambda \rightarrow Y$ connecting f_0 to f_1 and a 2-parametric family of injective homomorphisms $\Phi_{s,t}: T\Lambda \rightarrow TY$ such that $\Phi_{0,t} = df_t$ for all $t \in [0, 1]$, $\Phi_{s,0} = df_0$ and $\Phi_{s,1} = df_1$ for all $s \in [0, 1]$, and such that $\Phi_{1,t}$ is a Lagrangian homomorphism $T\Lambda \rightarrow \xi$ for all $t \in [0, 1]$. We will need the following simple lemma, see [11, 4, 18].

LEMMA 2.2. *Let $\Lambda \subset Y$ be a Legendrian submanifold and Λ^U its U -stabilization. Then if the Euler characteristic $\chi(U)$ of U satisfies $\chi(U) = 0$, then Λ and Λ^U are formally Legendrian isotopic.*

2.2. Loose Legendrian submanifolds. Let Y be a contact $(2n - 1)$ -manifold, $n > 2$. We continue using the notation from Section 2.1. A Legendrian embedding $\Lambda \rightarrow Y$ of a connected manifold Λ (which we sometimes simply call a *Legendrian knot*) is called *loose* if it is isotopic to the stabilization of another Legendrian knot. We point out that looseness depends on the ambient manifold. A loose Legendrian embedding Λ into a contact manifold Y need not be loose in a smaller neighborhood Y' , $\Lambda \subset Y' \subset Y$.

Any Legendrian submanifold $\Lambda \subset Y$ can be made loose by stabilizing it in arbitrarily small neighborhood of a point. Moreover, it can be made loose even without changing its formal Legendrian isotopy class. Indeed, one can first stabilize it and then U -stabilize for some U with $\chi(U) = -1$.

The following h -principle for loose Legendrian knots in contact manifolds of dimension $2n - 1 > 3$ is proved in [18]:

THEOREM 2.3 ([18]). *Any two loose Legendrian embeddings which coincide outside a compact set and which can be connected by a formal compactly supported Legendrian isotopy can be connected by a genuine compactly supported Legendrian isotopy.*

REMARK 2.4. It is also shown in [18] that the U -stabilization of a Legendrian knot is loose for any non-empty U .

These results imply the following:

COROLLARY 2.5. *Any loose Legendrian knot Λ is a stabilization of some other loose Legendrian knot Λ' .*

Proof. Let $U \subset \text{int}(\mathbb{R}_+^{n-1})$ be a ball and $C \subset \text{int}(\mathbb{R}_+^{n-1})$ be a domain with Euler characteristic -1 disjoint from U . If Λ is loose then according to Lemma 2.2 the stabilization $\Lambda^{C \cup U} = (\Lambda^C)^U$ is formally Legendrian isotopic to Λ . Hence in view of Theorem 2.3 there is a genuine Legendrian isotopy connecting the stabilization $(\Lambda^C)^U$ of the loose Legendrian knot $\Lambda' = \Lambda^C$ to Λ . \square

3. LAGRANGIAN IMMERSIONS WITH A CONICAL SINGULAR POINT

In this section we establish an h -principle for maps which are self-transverse Lagrangian immersions with the minimal possible number of double points away from a single conical singularity. This result is a generalization and a corollary of the corresponding result for Lagrangian embeddings from [13] which we state as Theorem 3.5 below.

3.1. Legendrian isotopy and Lagrangian concordance. The following result about realizing a Legendrian isotopy as a Lagrangian embedding of a cylinder is proved in [12]. Let Y be a contact manifold with contact structure ξ given by the contact 1-form α , $\xi = \ker(\alpha)$. The symplectization of Y is the manifold $Y \times \mathbb{R}$ with symplectic form $d(e^s \alpha)$, where s is a coordinate along the \mathbb{R} -factor.

Let $f_t: \Lambda \rightarrow Y$, $t \in [-1, 1]$, be a Legendrian isotopy that is constant near its endpoints and which connects Legendrian embeddings $f_- = f_{-1}$ and $f_+ = f_1$. We extend f_t to all $t \in \mathbb{R}$ by setting $f_t = f_-$ for $t \leq -1$ and $f_t = f_+$ for $t \geq 1$.

LEMMA 3.1. *There exists a Lagrangian embedding*

$$G: \Lambda \times \mathbb{R} \rightarrow Y \times \mathbb{R},$$

given by the formula $G(x, t) = (\tilde{f}_t(x), h(x, t))$ and with the following properties:

- *there exists $T > 0$ such that $G(x, t) = (f_-(x), t)$ for $t < -T$ and $G(x, t) = (f_+(x), t)$ for $t > T$;*
- *\tilde{f}_t is a Legendrian embedding C^∞ -close to f_t .*

We will need the following modification of Lemma 3.1 for Lagrangian immersions. Let $f_t: \Lambda \rightarrow Y$, $t \in [-1, 1]$, be a self-transverse regular Legendrian homotopy constant near its endpoints that connects Legendrian embeddings f_- and f_+ . As in Lemma 3.1 we extend f_t to all $t \in \mathbb{R}$ as independent of t for $|t| \geq 1$.

LEMMA 3.2. *There exists a self-transverse Lagrangian immersion*

$$G: \Lambda \times \mathbb{R} \rightarrow Y \times \mathbb{R},$$

given by the formula $G(x, t) = (\tilde{f}_t(x), h(x, t))$ and with the following properties:

- *there exists $T > 0$ such that $G(x, t) = (f_-(x), t)$ for $t < -T$ and $G(x, t) = (f_+(x), t)$ for $t > T$;*
- *\tilde{f}_t is C^∞ -close to f_t ;*
- *the double points of G are in one-to-one index preserving correspondence with the double points of the regular homotopy f_t .*

Proof. The construction from [12] which proves Lemma 3.1 can be applied with some additional care near the self-intersection instances of the Legendrian regular homotopy to prove Lemma 3.2. Here, however, we will use a different argument.

Note that it is sufficient to consider the case when there is exactly one transverse self-intersection point $q \in Y$ of the regular homotopy f_t at the moment $t = 0$. It is also sufficient to construct the immersed Lagrangian cylinder that corresponds to the Legendrian regular homotopy restricted to some interval $[-\epsilon, \epsilon]$ for a small $\epsilon > 0$, because then one can apply Lemma 3.1 for the isotopy f_t restricted to the rest of \mathbb{R} .

There exist local coordinates $(x, y, z) \in \mathbb{R}^{n-1} \times \mathbb{R}^{n-1} \times \mathbb{R} = \mathbb{R}_{\text{st}}^{2n-1}$ in a neighborhood Ω of q such that the two intersecting branches B_0 and B_1 of f_0 at q are given by the inclusion of

$$B_0 = \{(x, y, z) : y = 0, z = 0, |x| < 1\}, \quad (3.1)$$

$$B_1 = \{(x, y, z) : x = 0, z = 0, |y| < 1\}. \quad (3.2)$$

into $\mathbb{R}_{\text{st}}^{2n-1}$.

Modifying the regular homotopy f_t slightly for t close to 0 we obtain a regular homotopy \tilde{f}_t without self-intersections for $t \in [0, \epsilon]$ and which, for $t \in [-\epsilon, 0]$, is supported in Ω and has the following special properties:

- $\tilde{f}_t \equiv \tilde{f}_0$ on B_1 and near ∂B_0 ;
- $\tilde{f}_{-\epsilon}|_{B_0}$ is given by the formula

$$\tilde{f}_{-\epsilon}(\xi) = (x(\xi), y(\xi), z(\xi)) = (\xi, 0, -\delta), \quad \xi \in \mathbb{R}^{n-1}, |\xi| \leq 1$$

and, using the same notation, $\tilde{f}_0|_{B_0}$ is given by the formula

$$\xi \mapsto \left(\xi, \frac{\partial \phi}{\partial \xi}, \phi(\xi) \right),$$

where $\phi(\xi) = \delta\theta(|\xi|)$ for a small positive constant $\delta \ll \epsilon$ and a C^∞ -function $\theta: [0, 1] \rightarrow [-1, 1]$, which is equal to -1 near 1, is equal to 1 near 0, and which has non-positive derivative.

The isotopy $\{\tilde{f}_t\}_{t \in [0, \epsilon]}$ can be lifted to a Lagrangian cylinder in the symplectization using Lemma 3.1. Consider a C^∞ -function $\sigma: B_0 \times [-\epsilon, 0] \rightarrow \mathbb{R}$ with the following properties (using coordinates $(\xi, \tau) \in \mathbb{R}^{n-1} \times [-\epsilon, 0]$, $|\xi| < 1$ in analogy with the above):

- $\sigma(\xi, \tau) = -e^\tau \delta$ near $(B_0 \times \{-\epsilon\}) \cup (\partial B_0 \times [-\epsilon, 0])$;
- $\sigma(\xi, \tau) = e^\tau \phi(\xi)$ near $B_0 \times 0$;
- the function $\sigma(0, \tau)$, $\tau \in [-\epsilon, 0]$ has a unique zero in $(-\epsilon, 0)$, which is moreover a regular value.

Now the required Lagrangian immersion $G|_{(B_0 \cup B_1) \times [-\epsilon, 0]}$ lifting $\{\tilde{f}_t\}_{t \in [-\epsilon, 0]}$ to the symplectization $\mathbb{R}_{\text{st}}^{2n-1} \times \mathbb{R}$ can be defined by the formula

$$\begin{aligned} G(\eta, t) &= (x(\eta, \tau), y(\eta, \tau), z(\eta, \tau), t(\eta, \tau)) \\ &= (0, \eta, 0, \tau), & \text{if } (0, \eta, 0) \in B_1, \\ G(\xi, \tau) &= (x(\xi, \tau), y(\xi, \tau), z(\xi, \tau), t(\xi, \tau)) \\ &= \left(\xi, e^{-\tau} \frac{\partial \sigma}{\partial \xi}(\xi, \tau), e^{-\tau} \frac{\partial \sigma}{\partial \tau}(\xi, \tau), \tau \right), & \text{if } (\xi, 0, 0) \in B_0. \end{aligned}$$

Note that $G|_{B_0}$ and $G|_{B_1}$ are Lagrangian embeddings with respect to the symplectic form $\omega = d(e^t(dz - ydx))$. This is obvious for $G|_{B_1}$, for $G|_{B_0}$ we calculate

$$G^*(d(e^t(dz - ydx))) = d\left(e^\tau d\left(e^{-\tau} \frac{\partial \sigma}{\partial \tau}\right) - \frac{\partial \sigma}{\partial \xi} d\xi\right) = d\left(-\frac{\partial \sigma}{\partial \tau} d\tau - \frac{\partial \sigma}{\partial \xi} d\xi\right) = 0.$$

We also have $G(\xi, -\epsilon) = (\xi, 0, -\delta, -\epsilon) = (\tilde{f}_{-\epsilon}, -\epsilon)$ and $G(\xi, 0) = \left(\xi, \frac{\partial \phi}{\partial \xi}, \phi, 0\right) = (\tilde{f}_0, 0)$. On the other hand, the last property of the function σ guarantees that $G(B_0)$ and $G(B_1)$ intersect transversely at a unique point and that the index of intersection is the same as the self-intersection index of the regular homotopy f_t . \square

3.2. Lagrangian immersions with a conical point. Let $S_{\text{st}}^{2n-1} = (S^{2n-1}, \xi_{\text{st}})$ be the sphere with the standard contact structure ξ_{st} defined by the restriction α_{st} to the unit sphere $S^{2n-1} \subset \mathbb{R}^{2n}$ of the Liouville form $\lambda_{\text{st}} := \frac{1}{2} \sum_{j=1}^n (x_j dy_j - y_j dx_j)$ in $\mathbb{R}_{\text{st}}^{2n}$.

For any integer m we denote by $(x, r) \in S^{m-1} \times (0, \infty)$ polar coordinates in $\mathbb{R}^m - \{0\}$, i.e. x is the radial projection of a point to the unit sphere, and r is its distance to the origin. The symplectic form ω_{st} in \mathbb{R}^{2n} has the form $d(r^2 \alpha_{\text{st}})$ in polar coordinates.

A map $h: \mathbb{R}^n \rightarrow \mathbb{R}_{\text{st}}^{2n}$ is called a *Lagrangian cone* if $h^{-1}(0) = 0$ and if it is given by the formula $h(x, r) = (\phi(x), cr^2)$ in polar coordinates, where $\phi: S^{n-1} \rightarrow S_{\text{st}}^{2n-1}$ is a Legendrian embedding and c is a positive constant.

Note that there exists a symplectomorphism

$$C: (S^{2n-1} \times \mathbb{R}, d(e^t \alpha_{\text{st}})) \rightarrow (\mathbb{R}^{2n} - \{0\}, \omega_{\text{st}}) \quad (3.3)$$

given by the formula $C(x, t) = \left(x, e^{\frac{t}{2}}\right)$ in polar coordinates. Under this symplectomorphism Lagrangian cones in $\mathbb{R}_{\text{st}}^{2n}$ correspond to cylindrical Lagrangian manifolds in the symplectization $(S^{2n-1} \times \mathbb{R}, d(e^t \alpha_{\text{st}}))$ of S_{st}^{2n-1} .

Let L be an n -dimensional manifold and X a $2n$ -dimensional symplectic manifold. A map $f: L \rightarrow X$ is called a *Lagrangian immersion with a conical point* at $p \in L$, if $f|_{L-\{p\}}$ is a Lagrangian immersion, and if in a neighborhood of p and in a Darboux chart around $f(p)$, the map is equivalent to a Lagrangian cone $h: \mathbb{R}^n \rightarrow \mathbb{R}_{\text{st}}^{2n}$ near the origin. The Legendrian embedding $\phi: S^{n-1} \rightarrow S_{\text{st}}^{2n-1}$ corresponding to this cone is called the *link* of the conical point.

We define a regular Lagrangian homotopy $f_t: L \rightarrow X$, $t \in [0, 1]$, of immersions with a conical point at p to be a homotopy which is fixed in some neighborhood of the singular point p and that is an ordinary regular Lagrangian homotopy when restricted to $L - \{p\}$. For self-transverse immersions f with a conical point at p , we define the self-intersection number $I(f)$ as $I(f|_{L-\{p\}})$. Then $I(f)$ is invariant under regular homotopies fixed near p . We will need the following result.

LEMMA 3.3. *Let $h: \mathbb{R}^n \rightarrow \mathbb{R}_{\text{st}}^{2n}$ be a Lagrangian cone over a Legendrian link $\phi: S^{n-1} \rightarrow S_{\text{st}}^{2n-1}$. Then there exists a Hamiltonian regular homotopy $h_t: \mathbb{R}^n \rightarrow \mathbb{R}_{\text{st}}^{2n}$, $t \in [0, 1]$, with*

$h_0 = h$, which is fixed near the singularity and outside of a ball $B_R \subset \mathbb{R}^n$ of some radius $R > 0$ centered at 0, and such that the following hold:

- h_1 coincides with the cone over a loose Legendrian knot $\tilde{\phi}$ near $\partial B_{\frac{R}{2}}$;
- the immersion $h_1^- := h_1|_{h_1^{-1}(B_{R/2})}$ has exactly one transverse self-intersection point;
- if $n = 2k$ then, for any ϕ , we can arrange that $I(h_1^-) = (-1)^{k-1}$, and if in addition ϕ is assumed to be a loose Legendrian knot, then we can arrange also that $I(h_1^-) = (-1)^k$.

REMARK 3.4. Note that by scaling we can make the radius of the ball R arbitrarily small.

Proof. Let $\phi: S^{n-1} \rightarrow S_{\text{st}}^{2n-1}$ be the Legendrian link of the conical point. Let us denote by ϕ_1 its stabilization. Note that ϕ_1 is a loose knot. If ϕ is itself loose then according to Corollary 2.5 there exists another loose Legendrian knot ϕ_{-1} such that ϕ is the stabilization of ϕ_{-1} . We will call ϕ_{-1} the *destabilization* of ϕ . According to Lemma 2.1 the embeddings $\phi_0 := \phi$ and ϕ_1 can be included into a regular Legendrian homotopy $\phi_t: S^{n-1} \rightarrow S_{\text{st}}^{2n-1}$, $t \in [0, 1]$, such that there is exactly one self-intersection point for $t \in (0, 1)$, and when $n = 2k$ we have $I(\{\phi_t\}_{t \in [0, 1]}) = (-1)^{k-1}$. Similarly, if ϕ is loose then ϕ_{-1} and $\phi_0 = \phi$ can be included into a regular Legendrian homotopy $\phi_t: S^{n-1} \rightarrow S_{\text{st}}^{2n-1}$, $t \in [-1, 0]$, with one transverse self-intersection point which for an even $n = 2k$ has index $(-1)^{k-1}$.

Next, we use Lemma 3.2 to lift the Legendrian regular homotopy $\{\phi_t\}_{t \in [0, 1]}$ and its inverse $\{\phi_{1-t}\}_{t \in [0, 1]}$, and in the loose case $\{\phi_t\}_{t \in [-1, 0]}$ and its inverse $\{\phi_{-t}\}_{t \in [0, 1]}$, respectively, to Lagrangian immersions

$$G_1, G_2, G_3, G_4: S^{n-1} \times \mathbb{R} \rightarrow (S^{2n-1} \times \mathbb{R}, d(e^t \alpha_{\text{st}}))$$

with the following properties

- for a sufficiently large positive t we have

$$\begin{aligned} G_1(x, -t) &= (\phi(x), -t), & G_1(x, t) &= (\phi_1(x), t), \\ G_2(x, -t) &= (\phi_1(x), -t), & G_2(x, t) &= (\phi(x), t), \\ G_3(x, -t) &= (\phi_{-1}(x), -t), & G_3(x, t) &= (\phi(x), t), \\ G_4(x, -t) &= (\phi(x), -t), & G_4(x, t) &= (\phi_{-1}(x), t); \end{aligned}$$

- each of these Lagrangian immersions has exactly 1 transverse self-intersection point;
- if $n = 2k$ then

$$I(G_1) = I(G_3) = (-1)^{k-1}, \quad I(G_2) = I(G_4) = (-1)^k.$$

Composing these immersions with the symplectomorphism C^{-1} from (3.3) (and compactifying with a conical point) we get Lagrangian immersions

$$H_1, H_2, H_3, H_4: \mathbb{R}^n \rightarrow \mathbb{R}_{\text{st}}^{2n}$$

with a conical singularity at the origin. With appropriate rescaling we can glue together the immersions H_1 and H_2 to get an immersion $H_{12}: \mathbb{R}^n \rightarrow \mathbb{R}_{\text{st}}^{2n}$, and in the loose case glue H_4 and H_3 to get an immersion $H_{43}: \mathbb{R}^n \rightarrow \mathbb{R}_{\text{st}}^{2n}$, such that both are immersions with a conical singularity with Legendrian link ϕ , and both coincide with the cone over ϕ outside of the a ball B_R of some radius $R > 0$. In addition, near $\partial B_{R/2}$ the immersion H_{12} coincides with the cone over ϕ_1 and H_{43} coincides with the cone over ϕ_{-1} . Both immersions are homotopic to the Lagrangian cone h over ϕ via a Hamiltonian regular homotopy fixed outside B_R and near the singularity. A regular Hamiltonian homotopy connecting h with H_{12} is the required homotopy h_t in the general case. In the case of loose ϕ and even $n = 2k$, we can construct a Lagrangian immersion with the index of its unique self-intersection point equal to $(-1)^k$, by taking h_t to be a regular Hamiltonian homotopy connecting h with H_{43} . \square

3.3. The h -principle for self-transverse Lagrangian immersions with a conical singularity and minimal self-intersection. Let X be a symplectic $2n$ -manifold of dimension $2n > 4$. The following h -principle for Lagrangian *embeddings* into X with a conical point is proved in [13].

THEOREM 3.5. *Let $f_0: L \rightarrow X$ be a Lagrangian immersion with a conical point $p \in L$ into a simply connected symplectic manifold X of dimension $2n > 4$. If $n = 3$ we further assume that $X \setminus f_0(L)$ has infinite Gromov width, i.e. admits a symplectic embedding of an arbitrarily large ball. If the Legendrian link of f_0 at p is loose and if $I(f_0) = 0$, then there exists a Hamiltonian regular homotopy $f_t: L \rightarrow X$, $t \in [0, 1]$, that is fixed in a neighborhood of p and that connects f_0 to a Lagrangian embedding f_1 with a conical point at p .*

As we shall see, Theorem 3.5 generalizes to self-transverse Lagrangian immersions with a conical point of non-zero Whitney index and with the minimal number of self-intersection points. In fact, Theorem 3.5 itself is the key ingredient in the proof of its generalization, which we state next.

THEOREM 3.6. *Let $f_0: L \rightarrow X$ be a Lagrangian immersion with a conical point $p \in L$ into a simply connected symplectic manifold X of dimension $2n > 4$. If $n = 3$ we further assume that $X \setminus f_0(L)$ has infinite Gromov width. If the Legendrian link of f_0 at p is loose, then there exists a Hamiltonian regular homotopy $f_t: L \rightarrow X$, $t \in [0, 1]$, that is fixed in a neighborhood of p and that connects f_0 to a self-transverse Lagrangian immersion f_1 with a conical point at p and with $\text{SI}(f_1) = |I(f_0)|$.*

Proof. We argue by induction on $d = |I(f_0)|$, using Theorem 3.5 as the base of the induction for $d = 0$. Suppose the theorem holds for $|I(f_0)| = d - 1$. Let $\phi: S^{n-1} \rightarrow$

S_{st}^{2n-1} be the loose Legendrian link of the conical point and consider the immersion $f_0: L \rightarrow X$. By definition, in some local Darboux neighborhood near the singular point it is equivalent to a Lagrangian cone over ϕ in a ball B_ϵ , $\epsilon > 0$. We use Lemma 3.3 to construct a Hamiltonian regular homotopy supported in B_ϵ from f_0 to a new immersion \widehat{f}_0 that coincides with the cone over a loose knot $\widetilde{\phi}$ near $\partial B_{\epsilon/2}$ and which has exactly one transverse self-intersection point in $B_{\epsilon/2}$. If $n = 2k$, we arrange the index to be of the same sign as $I(f_0)$.

Let \widetilde{f}_0 be a Lagrangian immersion obtained from \widehat{f}_0 by modifying it to the cone over $\widetilde{\phi}$ in $B_{\epsilon/2}$. We note that $|I(\widetilde{f}_0)| = d - 1$, and by the induction hypothesis we find a Hamiltonian regular homotopy \widetilde{f}_t , $t \in [0, 1]$, fixed near the singularity and such that $\text{SI}(\widetilde{f}_1) = |I(\widetilde{f}_0)| = d - 1$. Note that the regular homotopy \widetilde{f}_t is fixed in B_σ for $\sigma \ll \frac{\epsilon}{2}$, but not necessarily fixed in $B_{\epsilon/2}$.

The required regular homotopy f_t is then obtained by deforming f_0 into \widehat{f}_0 , then scaling it inside $B_{\epsilon/2}$ to make it coincide with a cone in $B_{\epsilon/2} - B_\sigma$, and finally deforming it outside B_σ , keeping it fixed in B_σ , using the Hamiltonian regular homotopy \widetilde{f}_t . \square

4. PROOFS OF THE MAIN RESULTS

4.1. Proof of Theorem 1.1. Any simple (i.e. not double) point $p \in f_0(L)$ can be viewed as a conical point over a trivial Legendrian knot. Hence, we can apply Lemma 3.3 to find a Hamiltonian regular homotopy supported in a Darboux neighborhood of p , symplectomorphic to B_ϵ , such that the resulting Lagrangian immersion \widetilde{f}_0 coincides with a Lagrangian cone over a loose knot near $\partial B_{\epsilon/2}$ and has exactly one transverse self-intersection point in $B_{\epsilon/2}$, with index equal to $(-1)^{k-1}$ if $n = 2k$. Note that in all cases one then has $I(\widetilde{f}_0|_{L - \widetilde{f}_0^{-1}(B_{\epsilon/2})}) = I(f_0) + (-1)^k$. Hence, arguing as in the proof of Theorem 3.6, we replace $\widetilde{f}_0(L) \cap B_{\epsilon/2}$ by a cone over a loose Legendrian knot $\widetilde{\phi}$, and then using Theorem 3.6 we construct a Hamiltonian regular homotopy \widetilde{f}_t of the resulting immersion \widetilde{f}_0 , that is fixed in a neighborhood B_σ , $\sigma \ll \frac{\epsilon}{2}$, of the conical point, to an immersion with transverse self-intersections and with exactly $|I(f_0) + (-1)^k|$ double points. Finally the required regular homotopy f_t consists of first deforming \widetilde{f}_0 into \widehat{f}_0 , then scaling it inside $B_{\epsilon/2}$ in such a way that it becomes a cone in $B_{\epsilon/2} - B_\sigma$, and then deforming it outside B_σ using the Hamiltonian regular homotopy \widetilde{f}_t outside B_σ and keeping it fixed in B_σ . The Lagrangian immersion f_1 is self-transverse and has exactly

$$|I(f_0) + (-1)^{\frac{n}{2}}| + 1 = \begin{cases} |I(f_0)|, & \text{if } (-1)^{\frac{n}{2}} I(f_0) < 0, \\ |I(f_0) + 2|, & \text{if } (-1)^{\frac{n}{2}} I(f_0) \geq 0 \end{cases}$$

double points. \square

4.2. Proof of Corollary 1.3. Using Gromov's h -principle for Lagrangian immersions, see [16], we find an exact Lagrangian immersion $f: L \rightarrow \mathbb{R}_{\text{st}}^{2n}$ (in the given Lagrangian homotopy class σ in cases (1) and (3)). Part (1) then follows from the corresponding case of Theorem 1.1. If n is even then $I(f_0) = (-1)^{\frac{n}{2}} \frac{\chi(L)}{2}$, see e.g. [7], [Proposition 3.2] and also part (3) follows from the corresponding case of Theorem 1.1. To complete the proof of part (2), we observe that if $n = 3$ then both smooth regular homotopy classes of immersions $f: L \rightarrow \mathbb{R}_{\text{st}}^{2n}$, corresponding to $I(f) = 0$ and $I(f) = 1$, can be realized by a Lagrangian immersion. \square

4.3. Further results on $s(\sigma, L)$. If n is even then the smooth regular homotopy class of a Lagrangian immersion $f: L \rightarrow \mathbb{R}_{\text{st}}^{2n}$ is determined by $\chi(L)$, and thus Corollary 1.3 together with Gromov's non-existence result gives complete information on $s(L)$ if $\chi(L) \leq 0$. If $\chi(L) > 0$, $s(L)$ depends on more intricate, differential topological, properties of L , see [9, 10] (Theorem 1.4 gives information on the homotopy type of L in this case). For odd n , $s(L)$ (and $s(L, \sigma)$) is determined by which of the two smooth regular homotopy classes contain Lagrangian (or equivalently totally real) immersions. The following result gives a partial answer. Recall that $\chi_2(L) = \sum_{j=0}^{\frac{n-1}{2}} \text{rank}(H_j(L)) \pmod{2}$.

THEOREM 4.1. *If L is an n -dimensional orientable closed manifold with $TL \otimes \mathbb{C}$ trivial, n odd and $n \geq 3$, then the following hold:*

- (1) *If $n = 3$ then both regular homotopy classes contain exact Lagrangian immersions.*
- (2) *If $n \neq 2^q - 1$, $q \neq 2, 3$ then only one regular homotopy class contain Lagrangian immersions.*
- (3) *If $n = 4k + 1$ and $4k$ is not a power of two, or if the Stiefel-Whitney number $w_2(L)w_{n-2}(L)$ vanishes, then only the regular homotopy class with Whitney index $I_f = \chi_2(L)$ contains Lagrangian immersions.*
- (4) *If V is almost parallelizable then only the regular homotopy class with $I_f = \chi_2(V)$ contain Lagrangian immersions.*

Proof. Cases (1), (2), and (4) are proved in [2] and case (3) is a consequence of [1]. \square

4.4. Proof of Theorem 1.4. We control the homotopy type of exact immersions $f: L \rightarrow \mathbb{R}_{\text{st}}^{2n}$, n even, with exactly $\frac{1}{2}\chi(V) > 0$ double points using a straightforward generalization of [10, Lemma 2.2]. We refer to [10, Section 2] for background and notation for the parts of (lifted) Legendrian homology that will be used below.

Consider the Legendrian lift $\tilde{f}: L \rightarrow \mathbb{R}_{\text{st}}^{2n} \times \mathbb{R}$. The Reeb chords of \tilde{f} correspond to the double points of f , and the grading of all Reeb chords must be even since the sum of grading signs over Reeb chords equals $(-1)^{\frac{n}{2}}\chi(L)/2$, see [7]. Since no chord has odd grading it follows that the Legendrian algebra admits an augmentation, and since L is spin we can use arbitrary coefficients in the Legendrian algebra.

Using the duality sequence for linearized Legendrian homology [6] over \mathbb{Q} we find that all odd dimensional homology of L vanishes. In particular, the Maslov class vanishes and the linearized Legendrian homology admits an integer grading. The duality sequence with coefficients \mathbb{Z}_p for arbitrary prime p then implies that L has only even dimensional homology over \mathbb{Z} .

We next claim that $\pi_1(L) = 1$. To this end, consider a connected covering space $\pi: \tilde{L} \rightarrow L$ of L . Then the “lifted linearized Legendrian homology complex” $\tilde{C}^{\text{lin}}(\tilde{L}, \pi; k)$ with coefficients in the field k , see [10, Section 2], has the form

$$\tilde{S} \oplus \tilde{C}_{\text{Morse}} \oplus \tilde{L},$$

where the elements in \tilde{S} have grading $-1 + 2j$, $0 \leq j \leq \frac{n}{2}$ and the elements in \tilde{L} grading $2j + 1$, $0 \leq j \leq \frac{n}{2}$. Since $f(L)$ is Hamiltonian displaceable, the total homology of the complex vanishes. Since \tilde{L} is connected, this in turn implies $\text{rank}(\tilde{S}_{-1}) = 1$, where \tilde{S}_r is the degree r summand of \tilde{S} . Thus the covering π has degree 1, and since the covering was arbitrary, that implies that $\pi_1(V) = 1$, as required.

Finally, if $\dim L > 4$ then we can use the Whitney trick to cancel homologically inessential handles. \square

APPENDIX A. EXPLICIT CONSTRUCTIONS

In this section we consider explicit constructions of Lagrangian immersions and Lagrangian regular homotopies. In Sections A.1 and A.2 we introduce notation and some background material, which are necessary for the construction of a concrete Lagrangian immersion of $P = (S^1 \times S^{n-1}) \# (S^1 \times S^{n-1})$ into $\mathbb{R}_{\text{st}}^{2n}$. That construction, which is broken down into five stages, is given in the subsequent sections. The construction generalizes to dimensions $n \geq 3$ Sauvaget’s construction from [20] for $n = 2$.

A.1. Symplectization coordinates. Consider $\mathbb{R}_{\text{st}}^{2n}$ with coordinates $(x_1, y_1, \dots, x_n, y_n)$ and standard exact symplectic form $\omega = -d\beta$ with primitive $\beta = \sum_{j=1}^n y_j dx_j$. Let $\mathbb{C}^{n-1} \subset \mathbb{R}_{\text{st}}^{2n}$ denote the subspace given by the equation $(x_n, y_n) = (0, 0)$ and let $\beta_0 = \beta|_{\mathbb{C}^{n-1}}$. Consider the contact manifold $\mathbb{C}^{n-1} \times \mathbb{R}$ with contact 1-form $\alpha = dz - \beta_0$, where z is a coordinate in the additional \mathbb{R} -factor, and with symplectization $\mathbb{R} \times \mathbb{R} \times \mathbb{C}^{n-1}$ with exact symplectic form $d(e^t \alpha)$, where t is the coordinate in the symplectization direction. Write $\xi = (x_1, \dots, x_{n-1})$ and $\eta = (y_1, \dots, y_{n-1})$, then (ξ, η, x_n, y_n) are coordinates on $\mathbb{R}_{\text{st}}^{2n}$. Consider the map $\Phi: \mathbb{R} \times \mathbb{R} \times \mathbb{C}^{n-1} \rightarrow \mathbb{C}^n$,

$$\Phi(t, z, \xi, \eta) = (\xi, e^t \eta, e^t, z).$$

Then

$$\Phi^*\left(-\sum_j y_j dx_j\right) = -e^t(\eta \cdot d\xi + z dt) = e^t(dz - \eta \cdot d\xi) - d(e^t z) = \alpha - d(e^t z),$$

and hence Φ is an exact symplectomorphism from the symplectization $\mathbb{R} \times \mathbb{R} \times \mathbb{C}^{n-1}$ to $\mathbb{C}_+^n = \{(\xi, \eta, x_n, y_n) : x_n > 0\}$.

If $\Lambda \subset \mathbb{R} \times \mathbb{C}^{n-1}$ is a Legendrian submanifold then $\mathbb{R} \times \Lambda$ is an exact Lagrangian submanifold of the symplectization $\mathbb{R} \times \mathbb{R} \times \mathbb{C}^{n-1}$. If

$$(t, z(\lambda), \xi(\lambda), \eta(\lambda)), \quad \lambda \in \Lambda, t \in \mathbb{R}$$

is a parameterization of $\mathbb{R} \times \Lambda$ then its image under Φ is parameterized by

$$(\xi(\lambda), e^t \eta(\lambda), e^t, z(\lambda)) \in \mathbb{C}_+^n.$$

Conversely, if L is a conical Lagrangian submanifold in \mathbb{C}_+^{n-1} parameterized by

$$(\xi(\lambda), s\eta(\lambda)), s, y_n(\lambda)), \quad \lambda \in \Lambda, s \in \mathbb{R}_+,$$

then the image of L under Φ^{-1} is the cylinder on a Legendrian submanifold $\Lambda \subset \mathbb{R} \times \mathbb{C}^{n-1}$ parameterized by $(z(\lambda), \xi(\lambda), \eta(\lambda))$, $\lambda \in \Lambda$, where $z(\lambda) = y_n(\lambda)$.

A.2. Exact Lagrangian immersions by front slices. Let M be an n -manifold and let $f: M \rightarrow \mathbb{C}^n$ be an exact Lagrangian immersion. After small perturbation, the following general position properties hold:

- (1) All self-intersections of $f: M \rightarrow \mathbb{C}^n$ are transverse double points.
- (2) The coordinate function $x_n \circ f: M \rightarrow \mathbb{R}$ is a Morse function.
- (3) If $p = f(p_0) = f(p_1)$ is a double point of f then $x_n(p)$ is a regular value of $x_n \circ f$.

Assume that (1) – (3) hold. For any regular value a , the level set $M^a = (x_n \circ f)^{-1}(a)$ is a smooth $(n-1)$ -manifold which is the boundary of the sublevel set $M^{\leq a} = f^{-1}((-\infty, a])$, and if $\pi_n: \mathbb{C}^n \rightarrow \mathbb{C}^{n-1}$ is the projection that projects out (x_n, y_n) then $f^a = \pi_n \circ f: M^a \rightarrow \mathbb{C}^{n-1}$ is an exact Lagrangian immersion.

Double points of f are also double points of f^a . In order to determine which double points of f^a correspond to actual double points of f , we must recover the y_n -coordinate from f^a . Write $(x, y) = (\xi, \eta, x_n, y_n)$, where $\xi = (x_1, \dots, x_{n-1})$, $\eta = (y_1, \dots, y_{n-1})$, and (ξ, η) are standard coordinates on \mathbb{C}^{n-1} as above. If c is a double point of f^a , $c = f^a(c_0) = f^a(c_1)$ for $c_0 \neq c_1 \in M^a$ and if γ is a path connecting c_0 to c_1 in M^a then

$$y_n(c_1) - y_n(c_0) = \frac{d}{dx_n} \left(\int_{\gamma} \eta d\xi \right),$$

where $\eta d\xi = \sum_{j=1}^{n-1} y_j dx_j$. In other words, the y_n -coordinate at a double point is the x_n -derivative of the action of any path connecting its endpoints. Using the exactness of f , this can be rephrased as follows, if z is a primitive of $f^*(\beta)$ and $z^a = z|_{M^a}$ then

$$y_n(c_1) - y_n(c_0) = \frac{d}{da} (z^a(c_1) - z^a(c_0)). \quad (\text{A.1})$$

In our constructions below we will depict the exact Lagrangian slices by drawing their fronts in $\mathbb{R}^{n-1} \times \mathbb{R}$. Before discussing slices, we give a brief general description of fronts. Let N be a closed manifold and consider an exact Lagrangian immersion $g: N \rightarrow \mathbb{C}^n$. Pick a primitive $\zeta: N \rightarrow \mathbb{R}$ of $g^*(\beta)$. Then the map $G = (g, \zeta): N \rightarrow \mathbb{C}^n \times \mathbb{R}$ is a Legendrian immersion, everywhere tangent to the contact plane field $\ker(d\zeta - \beta)$ on $\mathbb{C}^n \times \mathbb{R}$. The front of g is the projection $\pi_F \circ G: N \rightarrow \mathbb{R}^n \times \mathbb{R}$, where π_F projects out the

y -coordinate. For generic g , the front has singularities; the front determines the original Lagrangian immersion via the equations

$$y_j = \frac{\partial z}{\partial x_j},$$

which admit solutions that can be extended continuously over the singular set (caustic) of the front. For generic g , double points of g lie off of the caustic and correspond to smooth points on the front with the same x -coordinate and with parallel tangent planes. The Reeb vector field of the contact form $dz - y dx$ on $\mathbb{C}^n \times \mathbb{R}$ is simply the coordinate vector field ∂_z and thus double points of g correspond to Reeb chords of G .

Below, we will study the fronts of exact Lagrangian immersions $f^a: M^a \rightarrow \mathbb{C}^{n-1}$ that are slices of a given Lagrangian immersion $f: M \rightarrow \mathbb{C}^n$, and as mentioned above it will be of importance to recover the y_n -coordinate at the double points of f^a . We will call the Reeb chords of the Legendrian lift of an exact slice f^a *slice Reeb chords*. Thus a slice Reeb chord is a vertical chord that connects two points on the front with parallel tangent planes. In our pictures below, we indicate the difference of the y_n -coordinate at the endpoints by showing whether the vertical chord that connects them grows or shrinks, cf. Equation (A.1).

A.3. Overview: a construction in five pieces. The rest of the Appendix is devoted to the construction of a self-transverse exact Lagrangian immersion $P = (S^1 \times S^{n-1} \# S^1 \times S^{n-1}) \rightarrow \mathbb{C}^n$ with exactly one double point. The construction of the immersion is broken down into five stages, two of which are described in terms of fronts, and three of which are described via Lagrangian slices.

In what follows, we write $\xi = (x_1, \dots, x_{n-1})$ and $t = x_n$. Let $\eta = (y_1, \dots, y_{n-1})$ be dual to ξ and $\tau = y_n$ dual to t . If $f(\cdot, t)$ is a function that depends on t , we write $f'(\cdot, t)$ for the partial derivative $\frac{\partial f}{\partial t}$.

The construction will be decomposed into the following five pieces: a bottom piece (i) which will be drawn in Section A.4 as a sequence of fronts and which contains five critical slices (i.e. slices containing critical points of the Morse function $x_n: P \rightarrow \mathbb{R}$), three middle pieces (ii) – (iv), see Sections A.6, A.7, and A.8 all without critical slices where we will draw the corresponding Lagrangian immersions into \mathbb{C}^n , and finally a top piece (v), also drawn by fronts, that contains the unique double point, see Section A.9.

A.4. The first piece of the immersion. The first piece is constructed in four steps.

A.4.1. We pass the minimum of the t -coordinate and a standard $(n-1)$ -sphere appears as shown in Figure 1. It has one slice Reeb chord c_1 of length $\ell(c_1)$ and $\ell'(c_1) > 0$.

A.4.2. We introduce two Bott families \mathbf{c}_2 and \mathbf{c}_3 of slice Reeb chords of index 0 and 1, respectively. Both families are topologically $(n-2)$ -spheres, symmetric about the central slice Reeb chord. The lengths of the slice Reeb chords of the families are $\ell(\mathbf{c}_2) < \ell(\mathbf{c}_3)$,




FIGURE 1. A standard sphere is born.




FIGURE 2. New born Bott families of shrinking chords.

$\ell'(\mathbf{c}_2) < 0$, and $\ell'(\mathbf{c}_3) < 0$; see Figure 2. It will be important later that $\ell(\mathbf{c}_3)$ is not too small compared to $\ell(c_1)$. We introduce the following quantities corresponding to certain areas in the Lagrangian slice projections which appear later, but here related to the lengths of the slice Reeb chords:

$$\ell(\mathbf{c}_3) = C, \quad \ell(\mathbf{c}_2) = -B + C, \quad \ell(c_1) = A - B + C,$$

where $C > 0$ and $A > B > 0$. Then $C' < 0$, $-B' + C' < 0$, and $A' > B' - C'$. To be definite about $\ell(\mathbf{c}_3)$, we take $B = \frac{\mu_0}{5}C$ and $A = \frac{\mu_1}{2}C$, where μ_j are parameters such that $\mu_j \approx 1$. (Since the slice Reeb chords must change length with t , we cannot enforce $\mu_j = 1$ but we can take these parameters to be approximately equal to 1.)

A.4.3. We Morse modify along the Bott family \mathbf{c}_2 of slice Reeb chords of minimal length twice as shown in Figure 3. It is straightforward to check that the manifold that results from these modifications is a punctured connected sum $(S^1 \times S^{n-1}) \# (S^1 \times S^{n-1})$, i.e., $P - D^n$. The slice sphere that appears right after the second Morse modification is depicted in Figure 4.

A.4.4. In the final step of the first piece, we Morsify the Bott family \mathbf{c}_2 of slice Reeb chords. In doing so, as we shall see below, it is natural to think about the quantities A ,




FIGURE 3. Morse modification along the Bott families of minima.




FIGURE 4. Final stage of the first piece.

B , and C as functions of $(v, t) \in S^{n-2} \times \mathbb{R}$, where we think of S^{n-2} as the unit sphere $\{\xi \in \mathbb{R}^{n-1} : |\xi| = 1\}$, which naturally parameterizes the Bott manifold. We Morsify \mathbf{c}_2 leaving one very short chord c_2^0 with $\ell(c_2^0) = \epsilon^2$ lying in direction $-v_0 \in S^{n-2}$ and one long chord c_2^{n-2} in direction $v_0 \in S^{n-2}$, with

$$\ell(c_2^{n-2}) = -B(v_0, t) + C = (1 - \frac{\mu_0}{5})C \approx \frac{4}{5}C, \quad (\text{A.2})$$

see Figure 4. Here, $\mu_0 \approx 1$ is as above, and C is constant in v since we keep the Bott symmetry of \mathbf{c}_3 . The superscripts on c_2^0 and c_2^{n-2} refer to the Morse indices of the slice Reeb chords, considered as positive function differences of the functions defining the two sheets of the front at their endpoints.

Let $\phi = \phi_+ - \phi_-$ be such a positive difference between local functions defining the sheets of the front. For future reference, we assume that the Morsification has only a small effect on the level surfaces of ϕ , which remain close to those of the Bott situation. In particular, the level curves of ϕ for values close to $\phi(c_2^{n-2})$ are everywhere transverse to the radial vectors along an S^{n-2} slightly outside the former Bott-manifold, see Figure 5.

A.5. A guide to reading pictures of Lagrangian slices. For the next three pieces of the immersion, we will not use the front representation as above, but will instead draw




FIGURE 5. Level surfaces after Morsification.

a family of exact Lagrangian slices. Before reaching the details we describe how to read the pictures.

Let $v \in S^{n-2} = \{\xi \in \mathbb{R}^{n-1} : |\xi| = 1\}$ and let $r \in \mathbb{R}_{\geq 0}$. We construct our exact Lagrangian slice-spheres by drawing their slices in the 2-dimensional half-planes determined by v and its dual vector w , i.e. w is a vector in η -space \mathbb{R}^{n-1} dual to v which lies in ξ -space. These slices are curves $\gamma_v : [0, 1] \rightarrow \{rv + \rho w : r \geq 0, \rho \in \mathbb{R}\}$ that begin and end at a central point (the location of the slice Reeb chord of maximal length of the corresponding front) at $(0, 0) = 0 \in \mathbb{C}^{n-1}$. In order for the curves γ_v to close up and form a sphere, the integral $\int_{\gamma_v} \eta d\xi$ must be independent of $v \in S^{n-2}$.

We will draw the immersed curves γ_v with over/under information recording the value of the $\tau = y_n$ -coordinate at double points. It is also important to keep track of the values of z at crossings, where z is a primitive function of the exact Lagrangian slice. We write $(\Delta\tau)_j$ and $(\Delta z)_j$ for the differences in τ -coordinate and z -coordinate, respectively, between the upper and the lower strands at the crossing labeled j in figures. Note that with these conventions $(\Delta z)_j \geq 0$.

We next need a description of the actual double points of the slice immersions f^a in terms of the curves γ_v . Note first that any double point of f^a corresponds to a double point of some curve γ_v . Although we break the S^{n-2} symmetry we stay fairly close to a symmetric situation. In particular, the double points of the curves γ_v (with the exception of the central double point) will come in S^{n-2} -families and the z -differences $(\Delta z)_j$ then give functions

$$(\Delta z)_j : S^{n-2} \rightarrow \mathbb{R}_{\geq 0}.$$

Since double points correspond to parallel tangent planes on the front we find that the double points of f^a are exactly the critical points of these functions.

A.6. The second piece of the immersion. We will represent the second piece of the immersion in three steps.




FIGURE 6. Initial Lagrangian slice.




FIGURE 7. Applying a finger move.

A.6.1. In the initial step the curves correspond to the front in Figure 4, which is depicted in the slice model in Figure 6. A , B , and C denote the (positively oriented) areas indicated; each is a function of v and t . We then have

$$\begin{aligned} (\Delta z)_1(t) &= A - B + C, & (\Delta \tau)_1 &= (A - B + C)' > 0, \\ (\Delta z)_2(v, t) &= -B + C, & (\Delta \tau)_2 &= (-B + C)' > 0, \\ (\Delta z)_3(v, t) &= -C, & (\Delta \tau)_3 &= (-C)' > 0. \end{aligned}$$

In particular $A(-v_0, t) > B(-v_0, t) = C(-v_0, t) - \epsilon^2$ and, for fixed t , the functions $A(v, t)$ and $B(v, t)$ have maxima at $-v_0$ and minima at v_0 , whereas the function $C(v, t)$ is constant in v .

A.6.2. We apply a finger move across the area $B(v, t)$, splitting it into two pieces $B_0(v, t)$ and $D(v, t)$ as shown in Figure 7, where $B_0(v, t)$ is constant in v :

$$B_0(v, t) = B_0(v_0, t) = D(v_0, t).$$

Consequently, $D(v, t)$ equals $B(v, t) - \text{const}$.

The crossing conditions then read:

$$\begin{aligned} (\Delta z)_1 &= A - B_0 + C - D, & (\Delta \tau)_1 &= (A - B_0 + C - D)' > 0, \\ (\Delta z)_2 &= -B_0 + C - D, & (\Delta \tau)_2 &= (-B_0 + C - D)' > 0, \\ (\Delta z)_3 &= -C, & (\Delta \tau)_3 &= (-C)' > 0, \\ (\Delta z)_4 &= -D, & (\Delta \tau)_4 &= (-D)' > 0. \end{aligned}$$




FIGURE 8. Final step of lower middle piece. The newly introduced area $E(v, t)$ is everywhere small at this stage.

A.6.3. We continue the finger move and introduce a new small area $E(v, t)$, as shown in Figure 8. We have

$$(\Delta z)_1 = A - B_0 + C - D + E, \quad (\Delta \tau)_1 = (A - B_0 + C - D + E)' > 0, \quad (\text{A.3})$$

$$(\Delta z)_2 = -B_0 + C - D + E, \quad (\Delta \tau)_2 = (-B_0 + C - D + E)' > 0, \quad (\text{A.4})$$

$$(\Delta z)_3 = -C, \quad (\Delta \tau)_3 = (-C)' > 0, \quad (\text{A.5})$$

$$(\Delta z)_4 = -D + E, \quad (\Delta \tau)_4 = (-D + E)' > 0, \quad (\text{A.6})$$

$$(\Delta z)_5 = -D, \quad (\Delta \tau)_5 = (-D)' > 0. \quad (\text{A.7})$$

A.7. The third piece of the immersion. The slices of the third piece will be drawn in the same way as the slices in Section A.6. We will distinguish small and large deformations, and use \approx_t for quantities that are almost constant in time, and $\approx_{(v,t)}$ for those that are almost constant in both $v \in S^{n-2}$ and in time. Recall that we wrote ϵ^2 for the height of the smallest Reeb chord, where $0 < \epsilon \ll 1$; by a small deformation we mean one smaller than ϵ^p for $p \gg 2$. For example $C(v, t) \approx_{(v,t)} \text{const}$ means that the area $C(v, t)$ is independent of v and that it has t -derivative of order ϵ^p , $p \gg 2$, with the sign of the derivative dictated by the crossing conditions.

A.7.1. In the initial phase of the third piece we change the area functions in such a way that the following conditions are met:

$$A(v, t) + E(v, t) \approx_{(v,t)} \text{const}, \quad B_0(v, t) \approx_{(v,t)} \text{const}, \quad C(v, t) \approx_{(v,t)} \text{const}, \quad D(v, t) \approx_t D(v, t'),$$

where t' is the starting time. Consider $v_0 \in S^{n-2}$ as above. At $-v_0$, the values of all area functions stay close to their initial values. At v_0 , $A(v, t)$ shrinks toward 0 and $E(v, t)$ grows correspondingly. We choose these functions so that they have exactly two critical points $\pm v_0$. Note that these deformations are compatible with (A.3)–(A.7). At the point when $A(v_0, t) = 0$ we find that the central slice Reeb chord which corresponds to a maximum of the function difference determined by the two sheets of the front and to the double point labeled 1 cancels with the chord c_2^{n-2} corresponding to the largest value of $(\Delta z)_2$. The curve γ_{v_0} at the cancelling moment is depicted in Figure 9. Note




FIGURE 9. The cancelling moment.




FIGURE 10. Level sets in the central region near the moment of cancellation.

that at this point the slice Reeb chord that corresponds to the double point labeled 4 satisfies the following:

$$(\Delta z)_4(v, t) < E(v_0, t) - D(v_0, t) \approx_t A(v_0, t') - D(v_0, t) < C(v, t), \quad (\text{A.8})$$

where we recall that $C(v, t) \approx_{(v, t)} \text{const}$ and that t' denotes the initial instance, before we start shrinking $A(v, t)$, see (i) *Step 4*. In particular, the Reeb chord of maximal length is the chord labeled 3 (denoted \mathbf{c}_3 earlier) of length $C(v, t)$.

A.7.2. In the central region where the Reeb chords cancel, the front of the Lagrangian slice consists of two function graphs. The level sets of the positive difference of these function graphs are shown in Figure 10.

Recall the consequence of the earlier Bott set up, see (i) *Step 4*, that the level sets are transverse to the radial vectors along an S^{n-2} surrounding the central region. Thus isotoping level sets keeping them fixed along the boundary, we see that there exists a Hamiltonian isotopy which is fixed outside the central region that deforms the Lagrangian so that the central region appears as shown in Figure 11. Here the slice Reeb chord c_2^0 is the new central Reeb chord and level sets are everywhere transverse to the radial vector




FIGURE 11. Placing the the minimal length Reeb chord in central position by an (almost) Hamiltonian deformation.




FIGURE 12. Appearance of the curves γ_v after c_2^0 became the new central Reeb chord.

field. Noting that double points in the central region of the radial slices γ_v correspond to tangencies of the level sets and the radial vector field, we find that after this Hamiltonian deformation, the curves γ_v are as shown in Figure 12. We take the slices of our exact Lagrangian to be approximately equal to the instances of this Hamiltonian deformation, deviating from it slightly in order to ensure that the crossing conditions hold.

We have the following crossing conditions:

$$\epsilon^2 \approx -\beta + C - D + E, \quad (\Delta\tau)_0 = (-\beta + C - D + E)' > 0, \quad (\text{A.9})$$

$$(\Delta z)_3 = -C, \quad (\Delta\tau)_3 = (-C)' > 0, \quad (\text{A.10})$$

$$(\Delta z)_4 = -D + E, \quad (\Delta\tau)_4 = (-D + E)' > 0, \quad (\text{A.11})$$

$$(\Delta z)_5 = -D, \quad (\Delta\tau)_5 = (-D)' > 0. \quad (\text{A.12})$$

For the deformation described above, the function $\beta(v, t)$ in Figure 12 has only two critical points, with maximum value at v_0 :

$$\beta(v_0, t) \approx A(-v_0, t) + B_0(v_0, t) > C(v, t), \quad (\text{A.13})$$

and minimum value at $-v_0$:

$$\beta(-v_0, t) \approx B_0(-v_0, t) < C(v, t). \quad (\text{A.14})$$




FIGURE 13. A rotation has been applied.

A.8. The fourth piece of the immersion. We describe the fourth piece in four steps.

A.8.1. We rotate the ends of the curves γ_v as shown in Figure 13. More precisely, recall that $(r, \rho) \mapsto rv + \rho w$ are coordinates on the half plane in which γ_v lies. Consider an interval $[0, R]$ and $r_0 \in [0, R]$ small, such that (i) all curves γ_v are of standard form in $\{(r, \rho): 0 \leq r \leq 2r_0\}$, and (ii) all curves γ_v are contained in $\{(r, \rho): 0 \leq r \leq \frac{1}{2}R\}$. Consider a Hamiltonian deformation ψ_t that is constant in $\{(r, \rho): 0 \leq r \leq r_0\}$, that is a $\frac{\pi}{2}$ rotation in the region $\{(r, \rho): 2r_0 \leq r \leq \frac{1}{2}R\}$, and that is again constant in $\{(r, \rho): \frac{3}{4}R \leq r\}$. The curves γ_v are then given by small deformations of the curves $\psi_t \circ \gamma_v$; the small deformations shrink the areas $C(v, t)$ and $D(v, t)$, and increase the area $\beta(v, t)$, in each case with very small derivative, in order to have the crossing conditions satisfied.

A.8.2. We shrink the area $C(v, t)$ for v in a certain subset of S^{n-2} . Recall from Equations (A.13), (A.14) that the minimum of the function $\beta(v, t)$ is $\beta(-v_0, t) \approx B_0(-v_0, t) = B(v_0, t) < C(v, t)$ whereas its maximum $\beta(v_0, t) \approx B(v_0, t) + A(-v_0, t) > C(v, t)$. Consider the subset $Q \subset S^{n-2}$ where



$$\beta(v, t) \geq C(v, t) - \frac{1}{2}\epsilon.$$

Over Q we shrink $C(v, t)$ and $\beta(v, t)$ in such a way that $C(v, t) - \beta(v, t) \approx \text{const}$ until $C(v, t) = \epsilon$. Outside a neighborhood of Q we keep $C(v, t)$ at its original size. Note that outside Q we have

$$E(v, t) = D(v, t) + \beta(v, t) - C(v, t) + \epsilon^2 \leq D(v, t) - \frac{1}{2}\epsilon + \epsilon^2 < D(v, t). \quad (\text{A.15})$$

A.8.3. We slide $C(v, t)$ out across $E(v, t)$ without creating double points. To see that this is possible we subdivide into three cases. First, around Q , $C(v, t)$ is small (size ϵ) compared to $E(v, t)$, which is of (order 1) finite size. In this case there are two double points created when $C(v, t)$ enters and leaves $E(v, t)$, see Figure 14. The crossing conditions at the entering crossings read

$$(\Delta\tau)_{\text{en}} = (\tilde{C}(v, t) - D(v, t) + E(v, t))' > 0,$$

FIGURE 14. Sliding a small area $C(v, t)$.FIGURE 15. Sliding a large area $C(v, t)$.

and those at the exiting crossings

$$(\Delta\tau)_{\text{ex}} = (\tilde{C}(v, t) - D(v, t))' > 0,$$

where $0 < \tilde{C}(v, t) < C(v, t) = \epsilon$. Here we let $E(v, t) \approx \text{const}$ and we let $D(v, t)$ shrink to keep the above derivative positive. As the total variation of $\tilde{C}(v, t)$ (i.e. the total amount of area transported in and out of $E(v, t)$) is 2ϵ and since $D(v, t) > 2\epsilon$, $D(v, t)$ is sufficiently large to keep the derivative positive at all times.

Second, well outside a neighborhood of Q , $C(v, t)$ is large and there are two double points throughout the deformation, see Figure 15, where the crossing conditions read

$$\begin{aligned} (\Delta\tau)_6 &= (\tilde{C}(v, t) - D(v, t) + \tilde{E}(v, t))' > 0, \\ (\Delta\tau)_7 &= (-D(v, t) + \tilde{E}(v, t))' > 0, \end{aligned}$$

where $\tilde{C}(v, t) \approx C(v, t) - \text{const}$ and where $0 \leq \tilde{E}(v, t) \leq E(v, t)$. Here $\tilde{E}'(v, t) < 0$ and we ensure that the crossing conditions hold by shrinking $D(v, t)$. Since the total variation of $\tilde{E}(v, t)$ is at most the area $E(v, t)$ and $D(v, t) > E(v, t)$ in this region, we find that the crossing condition is met at all times.

Third, in the region where we interpolate between small and large $C(v, t)$, $E(v, t)$ is already smaller than $D(v, t)$, and the amount of area transported through $E(v, t)$ is smaller than the corresponding amount described above; hence the necessary crossing conditions can be arranged by shrinking $D(v, t)$. In conclusion we can thus slide $C(v, t)$ out for all v .



FIGURE 16. Final shape of the curves γ_v .

FIGURE 17. Initial position for the fifth piece.

A.8.4. The final stage of the fourth piece is shown as a Lagrangian slice in Figure 16 and as the corresponding front in Figure 17. The latter serves as the starting point of the last piece.

A.9. The fifth piece of the immersion. For the fifth, final piece of the immersion, we return to the front representation. The initial front of the fifth piece is a sphere with seven Reeb chords, see Figure 17, three that shrink and four that grow. It is the double of a function graph with critical points as follows: (i) the largest maximum is positive and shrinking; (ii) there is another positive local maximum that increases; (iii) one positive index $n - 2$ point increases; (iv) there are two positive index 1 chords, one increasing and one decreasing; (v) and there are two index 0 chords, one positive increasing and one negative decreasing.

We pass the decreasing index 1 chord through a double point, making it increasing, see Figure 18. Then we cancel the two index $(0, 1)$ pairs and the index $(n - 1, n - 2)$ pair of increasing chords. This leaves us with a standard sphere, with a decreasing slice Reeb chord corresponding to a maximum, that we cap off with a Lagrangian disk via a standard Morse modification; see Figure 19. This completes the construction of the desired exact Lagrangian immersion $P \rightarrow \mathbb{C}^n$ with one double point.





FIGURE 18. Slices passing through the double point.

FIGURE 19. The final slice can be capped off with a maximum of x_n .

REFERENCES

- [1] Asadi-Golmankhaneh, M. and Eccles, P. *Double point self-intersection surfaces of immersions*. Geom. Topol. **4** (2000), 149–170. 4.3
- [2] Audin, M. *Fibrés normaux d’immersions en dimension double, points doubles d’immersions lagrangiennes et plongements totalement réels*. Comment. Math. Helv. **63** (1988), no. 4, 593–623. 4.3
- [3] Bourgeois, F., Ekholm, T., and Eliashberg, Y. *Effect of Legendrian surgery*. Geom. Topol. **16** (2012), no. 1, 301–389.
- [4] K. Cieliebak and Y. Eliashberg, *From Stein to Weinstein and back*
- [5] Damian, M. *Floer homology on the universal cover, a proof of Audin’s conjecture, and other constraints on Lagrangian submanifolds*. Comment. Math. Helv. **87** (2012) 433–462. 2.1, 2.1, 2.1 1, 1
- [6] Ekholm, T., Etnyre, J., and Sabloff, J. *A duality exact sequence for Legendrian contact homology*. Duke Math. J. **150** (2009), no. 1, 1–75 1, 4.4
- [7] T. Ekholm, J. Etnyre and M. Sullivan, *Non-isotopic Legendrian submanifolds in \mathbb{R}^{2n+1}* , J. Differential Geom. **71** (2005), 85–128. 4.2, 4.4
- [8] Ekholm, T., Etnyre, J., and Sullivan, M. *Orientations in Legendrian contact homology and exact Lagrangian immersions*. Internat. J. Math. **16** (2005), no. 5, 453–532. 1
- [9] Ekholm, T. and Smith, I. *Exact Lagrangian immersions with a single double point*. Preprint, arXiv:1111.5932. 1, 4.3
- [10] Ekholm, T. and Smith, I. *Exact Lagrangian immersions with one double point revisited*. Preprint, arXiv:1211.1715. 1, 1, 4.3, 4.4
- [11] Y. Eliashberg, *Topological characterization of Stein manifolds of dimension > 2* . Internat. J. Math. **1** (1990), 29–46. 2.1, 2.1

- [12] Y. Eliashberg and M. Gromov, *Lagrangian intersection theory: finite-dimensional approach*. In *Geometry of differential equations*, Amer. Math. Soc. Transl. Ser. 2, 186, (1998), 27–118. 3.1, 3.1
- [13] Eliashberg, Y. and Murphy, E. *Lagrangian caps* Preprint, 2013. (document), 1, 1, 2, 3, 3.3
- [14] K. Fukaya. *Application of Floer homology of Lagrangian submanifolds to symplectic topology*. In *Morse theoretic methods in nonlinear analysis and in symplectic topology*, 231–276, NATO Sci. Ser. II Math. Phys. Chem. 217, Springer (2006). 1
- [15] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono. *Lagrangian intersection Floer theory: anomaly and obstruction. Part I*. American Mathematical Society, International Press (2009). 1
- [16] Gromov, M. *Partial differential relations*. Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **9** Springer-Verlag, Berlin (1986). 4.2
- [17] Gromov, M. *Pseudoholomorphic curves in symplectic manifolds*. Invent. Math. **82** (1985), no. 2, 307–347.
- [18] E. Murphy, *Loose Legendrian Embeddings in High Dimensional Contact Manifolds*. Preprint, arXiv:1201.2245. 1, 2, 2.1, 2.1, 2.1, 2.2, 2.3, 2.4
- [19] L. Polterovich. *The surgery of Lagrange submanifolds*. Geom. Func. Anal. **2** (1991), 198–210. 1, 1
- [20] Sauvaget, D. *Curiosités lagrangiennes en dimension 4*. Ann. Inst. Fourier (Grenoble) **54** (2004) no. 6, 1997–2020. 1, A
- [21] C. Viterbo. *A new obstruction to embedding Lagrangian tori*. Invent. Math. **100** (1990), 301–320. 1
- [22] Whitney, H. *The self-intersections of a smooth n -manifold in $2n$ -space*. Ann. of Math. (2) **45**, (1944), 220–246. 1

TOBIAS EKHOLM, DEPARTMENT OF MATHEMATICS, UPPSALA UNIVERSITY, BOX 480, 751 06 UPPSALA, SWEDEN, AND INSTITUTE MITTAG-LEFFLER, AURAV 17, 182 60 DJURSHOLM, SWEDEN

YAKOV ELIASHBERG, DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, PALO ALTO, U.S.A.

EMMY MURPHY, DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, U.S.A.

IVAN SMITH, CENTRE FOR MATHEMATICAL SCIENCES, UNIVERSITY OF CAMBRIDGE, ENGLAND.